



**AG MIX**

**Herbicide**

A Sulfonylurea selective herbicide, for the post-emergence control of Wild Oats and Annual Phalaris and suppression of Brome grass, Barley grass and Annual ryegrass in Wheat.

**Active Ingredients:**  
Mesosulfuron Methyl 10g/l  
+ Iodosulfuron-methyl 2g/l  
+ Mefenpyr- Diethyl 30g/l  
**Product Type:**  
Oil Dispersion (OD)

**Agromec**

[www.agromec-international.com](http://www.agromec-international.com)



# AG MIX

# HERBICIDE



## Method of use:

Crop	Weeds	Leaf Stage	Rate of Use
Wheat	Brome grass	1 leaf to 3 Leaf, 1 tiller (Z11 to Z13-21)	330 ml/ Ha
	Annual Ryegrass	1 leaf to 3 Leaf, 1 tiller (Z11 to Z13-21)	330 ml/ Ha
	Barley grass	1 leaf to 4 Leaf, 2 tillers (Z11 to Z14-22)	330 ml/ Ha
	Wild oats	1 leaf to 3 Leaf, 1 tiller (Z11 to Z13-21)	330 ml/ Ha
	Annual phalaris, Phalaris paradoxa	1 leaf to 3 Leaf, 1 tiller (Z11 to Z13-21)	330 ml/ Ha

Withholding period:

Do not harvest for 8 weeks after application.